

# Effect of Nonlinear Thermal Radiation on Stagnation Flow of a Casson Fluid towards a Stretching Sheet

Ananth<sup>1</sup> P.A.Dinesh<sup>2</sup> V.Sugunamma<sup>3</sup> N.Sandeep<sup>4\*</sup>

<sup>1,3</sup> Department of Mathematics, S.V. University, Tirupati-517502, India.

<sup>2</sup> Department of Mathematics, M. S. Ramaiah Institute of Technology, Bangalore-560054.

<sup>3,4</sup> Fluid Dynamics Division, VIT University, Vellore-632014, India.

## Abstract

In this study, we analyzed the stagnation-point flow of a casson fluid towards a stretching sheet with nonlinear thermal radiation and induced magneticfield effects. The nonlinear partial differential equations are transformed into a set of nonlinear ordinary differential equations using self-suitable transformations, which are then solved numerically using Runge-Kutta based shooting technique. The effects of different non-dimensional parameters on velocity, induced magneticfield and temperature profiles along with the friction factor and local Nusselt number are discussed through graphs and tables. Comparisons of the present results with the existed literature were presented. Result indicate that the casson parameter have tendency to control the flow.

**Keywords:** Induced magneticfield, thermal radiation, casson fluid, stretching Sheet.

## 1. Introduction

Now a days the cooling of electronic devices is one of the important requirements in micro-electromechanical systems, nuclear reactors vehicle thermal management etc. The magneto hydrodynamics (MHD) deals with the motion of highly conducting fluids in the presence of magnetic field. The flow of the conducting fluid across the magneticfield causes the electric current, which has the capacity to change the influence of magneticfield and the flow behaviour of the fluid. It is important to study the flow characteristics of electrically conducting magneto-nanofluids. The magneto nanofluids have pivotal importance in the process of targeted drug release, asthma treatment, cancer therapy and in the construction of power generators. Flow due to stretching surface has many important applications in plastic and metal industries. Also, the glasses blowing, continuous casting of metals and spinning of fibers involve the concept of flow through stretching sheet. In all the above applications of flow through stretching sheet, the quality of the final product depends upon the rate of heat transfer at the stretching sheet.

The effect of non-linear thermal radiation on different types of fluid flows through various channels was studied by Hayat et al. (2015), Shehzad et al. (2014) and Mustafa et al. (2014). Cortell (2014) studied the nonlinear heat transfer effect on fluid flow over a stretching surface. Hayat et al. (2015a) discussed the MHD peristaltic flow of a nanofluid with slip effects and joule heating. The electro osmotic flow analysis of casson fluid flow past a micro slit channel was discussed by Chiu-On Ng (2013). Sulochana et al. (2015) discussed the influence of aligned magneticfield on the flow through vertical surface in porous medium with heat source. The flow characteristics on stagnation-point flow of casson fluid past a stretching surface in the presence of viscous dissipation effect was considered by Mustafa et al. (2012). An unsteady heat transfer characteristics on fluid flow past a permeable stretching surface with thermal radiation and non-uniform heat source/sink was considered by Pal (2011). Stagnation point flow of MHD Nanofluid flow past a non-isothermal stretching surface in presence of induced magneticfield effect was investigated by Pal and Mandal (2015). Jayachandra Babu et al. (2015) discussed the effect of radiation and viscous dissipation on stagnation-point flow of a micropolar fluid over a nonlinearly stretching surface.

Mohan Krishna et al. (2013, 2014) discussed the flow of dusty and nano fluids through different channels in the presence of transverse magneticfield. Aligned magneticfield, radiation and chemical reaction effects on dusty viscous flow over a flat plate were studied by Ramana Reddy et al. (2014). Sandeep et al. (2012, 2013) illustrated the radiation effects on the flows through porous media. An unsteady MHD free convection flow of a dissipative fluid past a vertical plate through porous media with constant heat flux was discussed by Sugunamma and Sandeep (2011). Radiation and inclined magnetic field effects on free convective flow past an impulsively moving vertical plate in a porous medium was studied by Sandeep and Sugunamma (2014). Raju et al. (2015) studied the influence of aligned magnetic field and radiation on the flow of ferrofluids over a flat plate with non-uniform heat source/sink. Sandeep and Sulochana (2015) presented dual solutions of radiative MHD nanofluid flow over an exponentially stretching sheet. Raju et al. (2015) illustrated the effects of radiation, inclined magnetic field and cross-diffusion effects on flow over a stretching surface. Stagnation point flow and heat transfer behavior of Cu-water nanofluid towards horizontal and exponentially stretching/shrinking cylinders was studied by Sulochana and Sandeep (2015). Sandeep and Sulochana (2015) analyzed the effects of MHD on the flow of a dusty nanofluid over a stretching surface with volume fraction of dust particles.

In this study, we analyzed the stagnation-point flow of a casson fluid towards a stretching sheet with

nonlinear thermal radiation and induced magneticfield effects. The nonlinear partial differential equations are transformed into a set of nonlinear ordinary differential equations using self-suitable transformations, which are then solved numerically using Runge-Kutta based shooting technique. The effects of different non-dimensional parameters on velocity, induced magneticfield and temperature profiles along with the friction factor and local Nusselt number are discussed through graphs and tables.

## 2. Mathematical Analysis

Consider a steady, incompressible, electrically conducting stagnation-point flow of a Casson fluid towards a stretching sheet in the presence of induced magneticfield. The stretching sheet is considered along the  $x$  – axis and  $y$  – axis is normal to it. It is assumed that the applied magneticfield is of uniform strength  $H_0$ . It is also assumed that induced magneticfield is applied in  $y$  – direction and the parallel component  $H_1$  approaches the value  $H_e = H_0$  in the free stream flow and normal component of the induced magneticfield  $H_2$  vanishes near the wall. Under the above assumptions the governing boundary layer equations are given by

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} = 0, \quad (1)$$

$$\frac{\partial H_1}{\partial x} + \frac{\partial H_2}{\partial y} = 0, \quad (2)$$

$$\rho \left( u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} \right) = \mu \left( 1 + \frac{1}{\beta} \right) \frac{\partial^2 u}{\partial y^2} + u_e(x) \frac{\partial u_e(x)}{\partial x} + \frac{\mu}{4\pi} \left( H_1 \frac{\partial H_1}{\partial x} + H_2 \frac{\partial H_1}{\partial y} - H_e \frac{\partial H_e}{\partial x} \right), \quad (3)$$

$$u \frac{\partial H_1}{\partial x} + v \frac{\partial H_1}{\partial y} = H_1 \frac{\partial u}{\partial x} + H_2 \frac{\partial u}{\partial y} + \alpha_1 \frac{\partial^2 H_1}{\partial y^2}, \quad (4)$$

$$u \frac{\partial T}{\partial x} + v \frac{\partial T}{\partial y} = \frac{k}{\rho c_p} \frac{\partial^2 T}{\partial y^2} + \frac{16\sigma^*}{3k^* \rho c_p} \frac{\partial}{\partial y} \left( T^3 \frac{\partial T}{\partial y} \right), \quad (5)$$

Subject to the boundary conditions

$$\left. \begin{aligned} u = u_w(x) = cx, v = 0, \frac{\partial H_1}{\partial y} = H_2 = 0, T = T_w \quad \text{at } y = 0, \\ u = u_e(x) = ax, H_1 = H_e(x) = H_0 x, T = T_\infty, \quad \text{as } y \rightarrow \infty, \end{aligned} \right\} \quad (6)$$

where  $u, v$  are the velocity components of the fluid in  $x, y$  directions,  $H_1, H_2$  are the magnetic components in  $x, y$  directions,  $\rho$  and  $\mu$  are the fluid density and dynamic viscosity respectively,  $\sigma$  is the electrical conductivity,  $\rho c_p$  is the specific heat capacitance,  $T$  is the fluid temperature,  $k$  is the effective thermal conductivity,  $\sigma^*$  and  $k^*$  are the Stefan-Boltzmann constant and the mean absorption coefficient, respectively,  $\alpha_1$  is the magnetic diffusivity of the nanofluid, which is given by  $\alpha_1 = 1 / 4\pi\sigma$ .

To convert the governing equations into set of nonlinear ordinary differential equations, we now introduce the following similarity transformation.

$$\left. \begin{aligned} u = cxf'(\eta), v = -v^{1/2}c^{1/2}f(\eta), \eta = v^{-1/2}c^{1/2}y, T = T_\infty(1 + (\theta_w - 1)\theta) \\ H_1 = H_0xg'(\eta), H_2 = -H_0v^{1/2}c^{-1/2}g(\eta), \theta(\eta) = (T - T_\infty) / (T_w - T_\infty), \end{aligned} \right\} \quad (7)$$

Substituting equations (6) and (7) into (1)-(5), equations (1) and (2) satisfies automatically. Now, the equations (3)-(5) will be transformed into the following nonlinear coupled ordinary differential equations:

$$\left( 1 + \frac{1}{\beta} \right) f''' - (f'^2 - ff'') + B(g'^2 - gg'' - 1) + \left( \frac{a}{c} \right)^2 = 0, \quad (8)$$

$$\lambda g''' + fg'' - f''g = 0, \quad (9)$$

$$\theta'' + R \left[ (1 + (\theta_w - 1)\theta)^3 \theta'' + 3(\theta_w - 1)\theta'^2 (1 + (\theta_w - 1)\theta)^2 \right] + \text{Pr} f\theta' = 0, \quad (10)$$

With the transformed boundary conditions

$$\left. \begin{aligned} f = 0, f' = 1, g = 0, g'' = 0, \theta = 1, & \quad \text{at } \eta = 0, \\ f' = a/c, g' = 1, \theta = 0, & \quad \text{as } \eta \rightarrow \infty, \end{aligned} \right\} \quad (11)$$

where  $\beta$  is the casson parameter,  $a, c$  are constants,  $B$  is the magnetic parameter,  $\lambda$  is the reciprocal magnetic Prandtl number,  $R$  is the radiation parameter,  $\theta_w$  is the temperature parameter,  $Pr$  the Prandtl number, which are represented below.

$$B = \frac{\mu H_0^2}{4\pi\rho c^2}, \lambda = \frac{1}{4\pi\sigma\nu}, R = \frac{16\sigma^* T_\infty^3}{3kk^*}, \theta_w = \frac{T_w}{T_\infty}, Pr = \frac{\nu}{\alpha}, \quad (12)$$

For physical quantities of engineering interest are the shear stress coefficient  $C_f$  (friction factor) and the local Nusselt number  $Nu_x$  are given by

$$Re_x^{1/2} C_f = f''(0), \quad (13)$$

$$Re_x^{-1/2} Nu_x = -\theta'(0), \quad (14)$$

Where  $Re_x$  is the local Reynolds number, given by

$$Re_x = \frac{u_w x}{\nu},$$

### 3. Results and Discussion

The system of nonlinear ordinary differential equations (8) – (10) with the boundary conditions (11) has been solved numerically using Runge-Kutta based shooting technique. Further the effects of various physical parameters like casson parameter  $\beta$ , magneticfield parameter  $B$ , temperature parameter  $\theta_w$ , radiation parameter  $R$  and stretching ratio  $a/c$  on velocity, induced magneticfield and temperature profiles. For numerical results we considered  $\theta_w = 1.1$ ,  $a/c = 0.5$ ,  $\beta = 0.5$ ,  $R = 1$ ,  $\lambda = 0.1$ . These values are kept as common in entire study except the varied values as displayed in the respective figures and tables.

Figs. 1 and 2 illustrate the effect of casson parameter  $\beta$  on velocity and induced magneticfield profiles. It is observed that an increase in the casson parameter declines the velocity as well as induced magneticfield profiles. This agrees the general physical behaviour of the non-Newtonian parameter. The effect of radiation parameter  $R$  on temperature profiles of the flow is shown in Fig. 3. It is found that an increase in the radiation parameter leads to an increase in the temperature profiles of the flow.

Figs. 4 and 5 depict the influence of magneticfield parameter on velocity and induced magneticfield profiles. It is evident that an increase in the magneticfield parameter causes to increase in the velocity and induced magneticfield profiles. This is due to the hike in the retardation force acts along the flow. Fig. 6 displays the effect of temperature parameter on the temperature profiles of the flow. It is clear that an increase in the temperature parameter enhances the temperature profiles of the flow. This may happen due to the enhanced thermal conductivity of the flow.

Fig. 7 and 8 reveal the influence of the stretching ratio parameter on velocity profiles respectively and induced magneticfield profiles. It is interesting to note that an increase in the value of  $a/c$  enhances the momentum boundary layer thickness and depreciates the induced magneticfield profiles. But we have noticed an opposite results to the above with an increase in the reciprocal magnetic Prandtl number, which are displayed in Figs. 9 and 10.

Finally the effects of various physical parameters on friction factor ( $C_f$ ) and local Nusselt number ( $Nu_x$ ) are shown in Table 1. It is evident that an increase in the casson parameter and reciprocal magnetic Prandtl number reduces the friction factor and heat transfer rate. A raise in the magneticfield parameter increases the friction factor and local Nusselt number. Stretching ratio parameter have tendency to increase the heat transfer rate and reduce the friction factor. Table 2 shows the validation of the present results with the existed literature under some special assumptions. We found an excellent agreement of the present results with the existed results.




Fig.1 Velocity field for different values of casson parameter




Fig.2 Induced magnetic field for different values of casson parameter




Fig.3 Temperature field for different values of radiation parameter




Fig.4 Velocity field for different values of magnetic parameter




Fig.5 Induced magnetic field for different values of magnetic parameter




Fig.6 Temperature field for different values of temperature parameter




Fig.7 Velocity field for different values of stretching ratio parameter




Fig.8 Induced magneticfield for different values of stretching ratio parameter




Fig.9 Velocity field for different values of reciprocal Prandtl number




Fig.10 Induced magneticfield for different values of reciprocal Prandtl number



**Table 1** Variation in friction factor ( $C_f$ ) and local Nusselt number ( $Nu_x$ )

$\beta$	$R$	$B$	$a/c$	$\lambda$	$Re_x^{1/2} C_f$	$Re_x^{-1/2} Nu_x$
0.5					-0.303296	1.209661
1.0					-0.372816	1.198239
1.5					-0.409458	1.192435
	1				-0.309131	1.208331
	2				-0.309131	0.946552
	3				-0.309131	0.800788
		0.5			-0.302736	1.209787
		1.0			-0.249572	1.219391
		1.5			-0.180341	1.232002
			0.4		-0.278463	1.203306
			0.5		-0.302736	1.209787
			0.6		-0.275401	1.215931
				0.5	-0.321107	1.207280
				1.0	-0.340453	1.203828
				1.5	-0.351822	1.201666

**Table 2** Comparison of the skin friction coefficient when  $R = 0$ .

$a/c$	Ali et al. (2011)	Present Results
0.1	-0.9694	-0.96935
0.2	-0.9181	-0.91820
0.5	-0.6673	-0.66737
2	2.0175	2.01749
3	4.7293	4.72934

#### 4. Conclusions

This study presents a stagnation-point flow of a casson fluid towards a stretching sheet with nonlinear thermal radiation and induced magneticfield effects. The nonlinear partial differential equations are transformed into a set of nonlinear ordinary differential equations using self-suitable transformations, which are then solved numerically using Runge-Kutta based shooting technique. The conclusions are as follows:

- Magneticfield parameter have tendency to enhance the heat transfer rate.
- Nonlinear thermal radiation enhances the thermal boundary layer thickness.
- Stretching ratio parameter helps to control the flow.
- Reciprocal magnetic Prandtl number have tendency to enhance the induced magneticfield profiles.
- Radiation parameter does not shown significant difference in the friction factor. But it helps to boost the temperature profiles of the flow.

#### References

- Ali F. M., R. Nazar, N. M. Arfin, I. Pop (2011). MHD stagnation-point flow and heat transfer towards stretching sheet with induced magnetic field, *Appl. Math. Mech. –Engl. Ed.*, 32(4), 409-418.
- Chiu-On Ng (2013). Combined pressure-driven and electroosmotic flow of casson fluid through a slit microchannel, *J. Non-Newtonian Fluid Mech.* 198, 1-9.
- Cortell R., (2014). Fluid flow and radiative nonlinear heat transfer over a stretching sheet, *Journal of King Saud University-Science* 26, 161-167.
- Hayat T., T. Muhammad, A. Alsaedi and M. S. Alhuthali, (2015). Magnetohydrodynamic three-dimensional flow of viscoelastic nanofluid in the presence of nonlinear thermal radiation, *Journal of Magnetism and Magnetic Materials* 385, 222-229
- Hayat T., Z. Nisar, B. Ahmad and H. Yasmin, (2015a). Simultaneous effects of slip and wall properties on MHD peristaltic motion of nanofluid with Joule heating, *Journal of Magnetism and Magnetic Materials* 395, 48-58.
- Jayachandra Babu, M., Radha Gupta, N.Sandeep. (2015). Effect of radiation and viscous dissipation on stagnation-point flow of a micropolar fluid over a nonlinearly stretching surface with suction/injection. *Journal of Basic and Applied Research International* 7(2), 73-82.
- Mustafa M, Tasawar Hayat, Ioan Pop, Awatif Hendi, (2012). Stagnation-point flow and heat transfer of a casson fluid towards a stretching sheet, *Z. Naturforsch.* 67, 70-76.

- Mustafa M., A. Mushtaq, T. Hayat and B. Ahmad, (2014). Nonlinear radiation heat transfer effects in the natural convective boundary layer flow of nanofluid past a vertical plate: A numerical study, *PLoS One* doi:10.1371/journal.pone.e0103946
- Mohan Krishna P, Sugunamma V and Sandeep N (2013). Magnetic field and chemical reaction effects on convective flow of a dusty viscous fluid, *Communications in Applied Sciences*. 1, 161-187.
- Mohan Krishna P, Sugunamma V, Sandeep N (2014). Radiation and magnetic field effects on unsteady natural convection flow of a nanofluid past an infinite vertical plate with heat source, *Chemical and Process Engineering Research*, Vol 25, pp39-52.
- Mohan Krishna P., N.Sandeep, V.Sugunamma. (2015) Effects of radiation and chemical reaction on MHD convective flow over a permeable stretching surface with suction and heat generation. *Walaliak Journal of Science and Technology*, 12(9), 831-847.
- Pal D. (2011). Combined effects of non-uniform heat source/sink and thermal radiation on heat transfer over an unsteady stretching permeable surface, *I. J. Communications in Nonlinear Science and Numeric Simula.* 16(4), 1890-1904.
- Pal, D., and Mandal, G. (2015). MHD convective stagnation-point flow of nanofluids over a non- isothermal stretching sheet with induced magnetic field, *Meccanica*. 1-13.
- Raju C.S. K., N. Sandeep, C. Sulochana, V. Sugunamma, (2015). Effects of aligned magnetic ` field and radiation on the flow of ferrofluids over a flat plate with non-uniform heat source/sink. *Internat. J. Sci. Eng.*, 8(2), 151-158.
- Raju C. S. K., N. Sandeep, C. Sulochana, V. Sugunamma, M. Jayachandra Babu, (2015). Radiation, inclined magnetic field and cross-diffusion effects on flow over a stretching surface. *JNNMS* 34, 169-180.
- Ramana Reddy, J.V., Sugunamma, V., Mohan Krishna, P., Sandeep, N. (2014). Aligned magnetic field, Radiation and chemical reaction effects on unsteady dusty viscous flow with heat generation/absorption, *Chemical and process eng. Research*, 27, 37- 53.
- Sandeep N, Reddy A.V.B, Sugunamma V, (2012). Effect of radiation and chemical reaction on transient MHD free convective flow over a vertical plate through porous media. *Chemical and process engineering research*. 2, 1-9.
- Sandeep, N., Sugunamma, V., Mohankrishna, P. (2013). Effects of radiation on an unsteady natural convective flow of a EG-Nimonic 80a nanofluid past an infinite vertical plate. *Advances in Physics Theories and Applications*. 23, 36-43.
- Sandeep N., V.Sugunamma (2014). Radiation and inclined magnetic field effects on unsteady MHD convective flow past an impulsively moving vertical plate in a porous medium. *Journal of Applied and Fluid Mechanics* 7(2), 275-286.
- Sandeep N, C. Sulochana, (2015). Dual solutions of radiative MHD nanofluid flow over an exponentially stretching sheet with heat generation/absorption. *Appl Nanosci* doi 10.1007/s13204-015-0420-z.
- Shehzad S. A., T. Hayat, A. Alsaedi and M. S.(2014). Alhuthali, Nonlinear thermal radiation in three-dimensional flow of Jeffery nanofluid: A model for solar energy, *Applied Mathematics and Computation* 248, 273-286.
- Sandeep N., C.Sulochana (2015). MHD flow of a dusty nanofluid over a stretching surface with volume fraction of dust particles. *Ain Shams Engineering Journal* (Press)
- Sugunamma, V. and Sandeep, N. (2011) .Unsteady hydromagnetic free convection flow of a dissipative and radiating fluid past a vertical plate through porous media with constant heat flux. *International journal of mathematics and computer applications research*. 1, 37-50.
- Sulochana C., N. Sandeep (2015), Stagnation point flow and heat transfer behavior of Cu- water nanofluid towards horizontal and exponentially stretching/shrinking cylinders. *Appl .Nanosci*; doi 10.1007/s13204-015-0451-5.
- Sulochana, M. , M.K.Kishore Kumar, N.Sandeep (2015). Influence of aligned magnetic field on the flow through vertical surface in porous medium with heat source. *Advances in Physics Theories and Applications* 42, 33-45.

The IISTE is a pioneer in the Open-Access hosting service and academic event management. The aim of the firm is Accelerating Global Knowledge Sharing.

More information about the firm can be found on the homepage:

<http://www.iiste.org>

## CALL FOR JOURNAL PAPERS

There are more than 30 peer-reviewed academic journals hosted under the hosting platform.

**Prospective authors of journals can find the submission instruction on the following page:** <http://www.iiste.org/journals/> All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Paper version of the journals is also available upon request of readers and authors.

## MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Academic conference: <http://www.iiste.org/conference/upcoming-conferences-call-for-paper/>

## IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

